In mathematics, the **logarithm** of a number is the exponent by which another fixed value, the **base**, must be raised to produce that number. For example, the logarithm of 1000 to base 10 is 3, because 1000 is 10 to the 3rd power: $1000 = 10^3 = 10 \times 10 \times 10$. More generally, if $x = b^y$, then y is the logarithm of x to base b, written $\log_b x$, so $\log_{10} 1000 = 3$. As a single-variable function, the logarithm to base b is the inverse of exponentiation with base b.

The logarithm base 10 is called the *decimal* or *common* logarithm and is commonly used in science and engineering. The *natural* logarithm has the number $e \approx 2.718$ as its base; its use is widespread in mathematics and physics because of its very simple derivative. The *binary* logarithm uses base 2 and is widely used in computer science, information theory, music theory, and photography. When the base is unambiguous from the context or irrelevant it is often omitted, and the logarithm is written log x.

Logarithms were introduced by John Napier in 1614 as a means of simplifying calculations.[1] They were rapidly adopted by navigators, scientists, engineers, surveyors, and others to perform high-accuracy computations more easily. Using logarithm tables, tedious multi-digit multiplication steps can be replaced by table look-ups and simpler addition. This is possible because the logarithm of a product is the sum of the logarithms of the factors:

$$\log_b(xy) = \log_b x + \log_b y,$$

provided that b, x and y are all positive and $b \neq 1$. The slide rule, also based on logarithms, allows quick calculations without tables, but at lower precision. The present-day notion of logarithms comes from Leonhard Euler, who connected them to the exponential function in the 18th century, and who also introduced the letter e as the base of natural logarithms.[2]

Logarithmic scales reduce wide-ranging quantities to smaller scopes. For example, the decibel (dB) is a unit used to express ratio as logarithms, mostly for signal power and amplitude (of which sound pressure is a common example). In chemistry, pH is a logarithmic measure for the acidity of an aqueous solution. Logarithms are commonplace in scientific formulae, and in measurements of the complexity of algorithms and of geometric objects called fractals. They help to describe frequency ratios of musical intervals, appear in formulas counting prime numbers or approximating factorials, inform some models in psychophysics, and can aid in forensic accounting.

The concept of logarithm as the inverse of exponentiation extends to other mathematical structures as well. However, in general settings, the logarithm tends to be a multi-valued function. For example, the complex logarithm is the multi-valued inverse of the complex exponential function. Similarly, the discrete logarithm is the multi-valued inverse of the exponential function in finite groups; it has uses in public-key cryptography.

Motivation

Addition, multiplication, and exponentiation are three of the most fundamental arithmetic operations. The inverse of addition is subtraction, and the inverse of multiplication is division. Similarly, a logarithm is the inverse operation of exponentiation. Exponentiation is when a number *b*, the *base*, is raised to a certain power *y*, the *exponent*, to give a value *x*; this is denoted

$b^y = x.$

For example, raising 2 to the power of 3 gives 8: $2^3 = 8$.

The logarithm of base *b* is the inverse operation, that provides the output *y* from the input *x*. That is, $y = \log_b x$ is equivalent to $x = b^y$ if *b* is a positive real number. (If *b* is not a positive real number, both exponentiation and logarithm can be defined but may take several values, which makes definitions much more complicated.)

One of the main historical motivations of introducing logarithms is the formula

$\log_b(xy) = \log_b x + \log_b y,$

by which tables of logarithms allow multiplication and division to be reduced to addition and subtraction, a great aid to calculations before the invention of computers.

Definition

Given a positive real number *b* such that $b \neq 1$, the *logarithm* of a positive real number *x* with respect to base *b*[nb 1] is the exponent by which *b* must be raised to yield *x*. In other words, the logarithm of *x* to base *b* is the unique real number *y* such that $b^y = x$.[3]

The logarithm is denoted "$\log_b x$" (pronounced as "the logarithm of *x* to base *b*", "the base-*b* logarithm of *x*", or most commonly "the log, base *b*, of *x*").

An equivalent and more succinct definition is that the function $\log_b$ is the inverse function to the function $x \mapsto b^x$.

Examples

- $\log_2 16 = 4$, since $2^4 = 2 \times 2 \times 2 \times 2 = 16$.
- Logarithms can also be negative: $\log_2 \frac{1}{2} = -1$ since $2^{-1} = \frac{1}{2^1} = \frac{1}{2}$.
- $\log_{10} 150$ is approximately 2.176, which lies between 2 and 3, just as 150 lies between $10^2 = 100$ and $10^3 = 1000$.
- For any base b, $\log_b b = 1$ and $\log_b 1 = 0$, since $b^1 = b$ and $b^0 = 1$, respectively.

Logarithmic identities

Several important formulas, sometimes called *logarithmic identities* or *logarithmic laws*, relate logarithms to one another.[4]

Product, quotient, power, and root

The logarithm of a product is the sum of the logarithms of the numbers being multiplied; the logarithm of the ratio of two numbers is the difference of the logarithms. The logarithm of the p-th power of a number is p times the logarithm of the number itself; the logarithm of a p-th root is the logarithm of the number divided by p. The following table lists these identities with examples. Each of the identities can be derived after substitution of the logarithm definitions $x = b^{\log_b x}$ or $y = b^{\log_b y}$ in the left hand sides. In the following formulas, x and y are positive real numbers and p is an integer greater than 1.

Product, quotient, power, and root identities of logarithms

Identity	Formula	Example

Product	$\log_b(xy) = \log_b x + \log_b y$	$\log_3 243 = \log_3(9 \cdot 27) = \log_3 9 + \log_3 27 = 2 + 3 = 5$
Quotient	$\log_b \frac{x}{y} = \log_b x - \log_b y$	$\log_2 16 = \log_2 \frac{64}{4} = \log_2 64 - \log_2 4 = 6 - 2 = 4$
Power	$\log_b\left(x^p\right) = p \log_b x$	$\log_2 64 = \log_2\left(2^6\right) = 6 \log_2 2 = 6$
Root	$\log_b \sqrt[p]{x} = \frac{\log_b x}{p}$	$\log_{10} \sqrt{1000} = \frac{1}{2} \log_{10} 1000 = \frac{3}{2} = 1.5$

Change of base

The logarithm $\log_b x$ can be computed from the logarithms of x and b with respect to an arbitrary base k using the following formula:[nb 2]

$$\log_b x = \frac{\log_k x}{\log_k b}.$$

Typical scientific calculators calculate the logarithms to bases 10 and e.[5] Logarithms with respect to any base b can be determined using either of these two logarithms by the previous formula:

$$\log_b x = \frac{\log_{10} x}{\log_{10} b} = \frac{\log_e x}{\log_e b}.$$

Given a number x and its logarithm $y = \log_b x$ to an unknown base b, the base is given by:

$$b = x^{\frac{1}{y}},$$

which can be seen from taking the defining equation $x = b^{\log_b x} = b^y$ to the power of $\frac{1}{y}$.

Particular bases

Among all choices for the base, three are particularly common. These are $b = 10$, $b = e$ (the irrational mathematical constant $e \approx 2.71828183$), and $b = 2$ (the binary logarithm). In mathematical analysis, the logarithm base e is widespread because of analytical properties

explained below. On the other hand, base 10 logarithms (the common logarithm) are easy to use for manual calculations in the decimal number system:[6]

$$\log_{10}(10\,x)=\log_{10} 10+\log_{10} x=1+\log_{10} x.$$

Thus, $\log_{10}(x)$ is related to the number of decimal digits of a positive integer x: The number of digits is the smallest integer strictly bigger than $\log_{10}(x)$.[7] For example, $\log_{10}(5986)$ is approximately 3.78 . The next integer above it is 4, which is the number of digits of 5986. Both the natural logarithm and the binary logarithm are used in information theory, corresponding to the use of nats or bits as the fundamental units of information, respectively.[8] Binary logarithms are also used in computer science, where the binary system is ubiquitous; in music theory, where a pitch ratio of two (the octave) is ubiquitous and the number of cents between any two pitches is a scaled version of the binary logarithm, or log 2 times 1200, of the pitch ratio (that is, 100 cents per semitone in conventional equal temperament), or equivalently the log base $2^{1/1200}$; and in photography, where rescaled base 2 logarithms are used to measure exposure values, light levels, exposure times, lens apertures, and film speeds in "stops".[9]

The abbreviation log x is often used when the intended base can be inferred based on the context or discipline, or when the base is indeterminate or immaterial. Common logarithms (base 10), historically used in logarithm tables and slide rules, are a basic tool for measurement and computation in many areas of science and engineering; in these contexts log x still often means the base ten logarithm.[10] In mathematics log x usually refers to the natural logarithm (base e).[11] In computer science and information theory, log often refers to binary logarithms (base 2).[12] The following table lists common notations for logarithms to these bases. The "ISO notation" column lists designations suggested by the International Organization for Standardization.[13]

Base b	Name for $\log_b x$	ISO notation	Other notations
2	binary logarithm	lb x [14]	ld x, log x, lg x,[15] $\log_2 x$
e	natural logarithm	ln x [nb 3]	log x, $\log_e x$
10	common logarithm	lg x	log x, $\log_{10} x$

b	logarithm to base *b*	$\log_b x$	

History

The history of logarithms in seventeenth-century Europe saw the discovery of a new function that extended the realm of analysis beyond the scope of algebraic methods. The method of logarithms was publicly propounded by John Napier in 1614, in a book titled *Mirifici Logarithmorum Canonis Descriptio* (*Description of the Wonderful Canon of Logarithms*).[19][20] Prior to Napier's invention, there had been other techniques of similar scopes, such as the prosthaphaeresis or the use of tables of progressions, extensively developed by Jost Bürgi around 1600.[21][22] Napier coined the term for logarithm in Middle Latin, *logarithmus*, literally meaning 'ratio-number', derived from the Greek *logos* 'proportion, ratio, word' + *arithmos* 'number'.

The common logarithm of a number is the index of that power of ten which equals the number.[23] Speaking of a number as requiring so many figures is a rough allusion to common logarithm, and was referred to by Archimedes as the "order of a number".[24] The first real logarithms were heuristic methods to turn multiplication into addition, thus facilitating rapid computation. Some of these methods used tables derived from trigonometric identities.[25] Such methods are called prosthaphaeresis.

Invention of the function now known as the natural logarithm began as an attempt to perform a quadrature of a rectangular hyperbola by Grégoire de Saint-Vincent, a Belgian Jesuit residing in Prague. Archimedes had written *The Quadrature of the Parabola* in the third century BC, but a quadrature for the hyperbola eluded all efforts until Saint-Vincent published his results in 1647. The relation that the logarithm provides between a geometric progression in its argument and an arithmetic progression of values, prompted A. A. de Sarasa to make the connection of Saint-Vincent's quadrature and the tradition of logarithms in prosthaphaeresis, leading to the term "hyperbolic logarithm", a synonym for natural logarithm. Soon the new function was appreciated by Christiaan Huygens, and James Gregory. The notation Log *y* was adopted by Gottfried Wilhelm Leibniz in 1675,[26] and the next year he connected it to the integral $\int \frac{dy}{y}$.

Before Euler developed his modern conception of complex natural logarithms, Roger Cotes had a nearly equivalent result when he showed in 1714 that[27]

$\log(\cos\theta + i\sin\theta) = i\theta.$

Logarithm tables, slide rules, and historical applications

By simplifying difficult calculations before calculators and computers became available, logarithms contributed to the advance of science, especially astronomy. They were critical to advances in surveying, celestial navigation, and other domains. Pierre-Simon Laplace called logarithms

> ... [a]n admirable artifice which, by reducing to a few days the labour of many months, doubles the life of the astronomer, and spares him the errors and disgust inseparable from long calculations.[28]

As the function $f(x) = b^x$ is the inverse function of $\log_b x$, it has been called an **antilogarithm**.[29] Nowadays, this function is more commonly called an exponential function.

Log tables

A key tool that enabled the practical use of logarithms was the *table of logarithms*.[30] The first such table was compiled by Henry Briggs in 1617, immediately after Napier's invention but with the innovation of using 10 as the base. Briggs' first table contained the common logarithms of all integers in the range from 1 to 1000, with a precision of 14 digits. Subsequently, tables with increasing scope were written. These tables listed the values of $\log_{10} x$ for any number x in a certain range, at a certain precision. Base-10 logarithms were universally used for computation, hence the name common logarithm, since numbers that differ by factors of 10 have logarithms that differ by integers. The common logarithm of x can be separated into an integer part and a fractional part, known as the characteristic and mantissa. Tables of logarithms need only include the mantissa, as the characteristic can be easily determined by counting digits from the decimal point.[31] The characteristic of $10 \cdot x$ is one plus the characteristic of x, and their mantissas are the same. Thus using a three-digit log table, the logarithm of 3542 is approximated by

$$\begin{aligned}\log_{10} 3542 &= \log_{10}(1000 \cdot 3.542) \\ &= 3 + \log_{10} 3.54\end{aligned}$$

Greater accuracy can be obtained by interpolation:

$$\log_{10} 3542 \approx 3 + \log_{10} 3.54 + 0.2\left(\log_{10} 3.55 - \log_{10} 3.54\right)$$

The value of 10^x can be determined by reverse look up in the same table, since the logarithm is a monotonic function.

Computations

The product and quotient of two positive numbers c and d were routinely calculated as the sum and difference of their logarithms. The product cd or quotient c/d came from looking up the antilogarithm of the sum or difference, via the same table:

$$cd = 10^{\log_{10} c} 10^{\log_{10} d} = 10^{\log_{10} c + \log_{10} d}$$

and

$$\frac{c}{d} = c\,d^{-1} = 10^{\log_{10} c - \log_{10} d}.$$

For manual calculations that demand any appreciable precision, performing the lookups of the two logarithms, calculating their sum or difference, and looking up the antilogarithm is much faster than performing the multiplication by earlier methods such as prosthaphaeresis, which relies on trigonometric identities.

Calculations of powers and roots are reduced to multiplications or divisions and lookups by

$$c^d = \left(10^{\log_{10} c}\right)^d = 10^{d \log_{10} c}$$

and

$$\sqrt[d]{c} = c^{\frac{1}{d}} = 10^{\frac{1}{d}\log_{10} c}.$$

Trigonometric calculations were facilitated by tables that contained the common logarithms of trigonometric functions.

Slide rules

Another critical application was the slide rule, a pair of logarithmically divided scales used for calculation. The non-sliding logarithmic scale, Gunter's rule, was invented shortly after Napier's invention. William Oughtred enhanced it to create the slide rule—a pair of logarithmic scales movable with respect to each other. Numbers are placed on sliding scales at distances proportional to the differences between their logarithms. Sliding

the upper scale appropriately amounts to mechanically adding logarithms, as illustrated here:



Schematic depiction of a slide rule. Starting from 2 on the lower scale, add the distance to 3 on the upper scale to reach the product 6. The slide rule works because it is marked such that the distance from 1 to x is proportional to the logarithm of x.

For example, adding the distance from 1 to 2 on the lower scale to the distance from 1 to 3 on the upper scale yields a product of 6, which is read off at the lower part. The slide rule was an essential calculating tool for engineers and scientists until the 1970s, because it allows, at the expense of precision, much faster computation than techniques based on tables.[32]

Analytic properties

A deeper study of logarithms requires the concept of a *function*. A function is a rule that, given one number, produces another number.[33] An example is the function producing the x-th power of b from any real number x, where the base b is a fixed number. This function is written as $f(x) = b^x$. When b is positive and unequal to 1, we show below that f is invertible when considered as a function from the reals to the positive reals.

Existence

Let b be a positive real number not equal to 1 and let $f(x) = b^x$.

It is a standard result in real analysis that any continuous strictly monotonic function is bijective between its domain and range. This fact follows from the intermediate value theorem.[34] Now, f is strictly increasing (for $b > 1$), or strictly decreasing (for $0 < b < 1$),[35] is continuous, has domain R, and has range $R_{>0}$. Therefore, f is a bijection from R to $R_{>0}$. In other words, for each positive real number y, there is exactly one real number x such that $b^x = y$.

We let $\log_b : R_{>0} \to R$ denote the inverse of f. That is, $\log_b y$ is the unique real number x such that $b^x = y$. This function is called the base-b *logarithm function* or *logarithmic function* (or just *logarithm*).

Characterization by the product formula

The function $\log_b x$ can also be essentially characterized by the product formula

$$\log_b(xy) = \log_b x + \log_b y.$$

More precisely, the logarithm to any base $b > 1$ is the only [increasing function] f from the positive reals to the reals satisfying $f(b) = 1$ and[36]

$$f(xy) = f(x) + f(y).$$

Graph of the logarithm function

As discussed above, the function $\log_b$ is the inverse to the exponential function $x \mapsto b^x$. Therefore, their [graphs] correspond to each other upon exchanging the x- and the y-coordinates (or upon reflection at the diagonal line $x = y$), as shown at the right: a point $(t, u = b^t)$ on the graph of f yields a point $(u, t = \log_b u)$ on the graph of the logarithm and vice versa. As a consequence, $\log_b(x)$ [diverges to infinity] (gets bigger than any given number) if x grows to infinity, provided that b is greater than one. In that case, $\log_b(x)$ is an [increasing function]. For $b < 1$, $\log_b(x)$ tends to minus infinity instead. When x approaches zero, $\log_b x$ goes to minus infinity for $b > 1$ (plus infinity for $b < 1$, respectively).

Derivative and antiderivative

Analytic properties of functions pass to their inverses.[34] Thus, as $f(x) = b^x$ is a continuous and [differentiable function], so is $\log_b y$. Roughly, a continuous function is differentiable if its graph has no sharp "corners". Moreover, as the [derivative] of $f(x)$ evaluates to $\ln(b)\, b^x$ by the properties of the [exponential function], the [chain rule] implies that the derivative of $\log_b x$ is given by[35][37]

$$\frac{d}{dx}\log_b x = \frac{1}{x \ln b}.$$

That is, the slope of the tangent touching the graph of the base-b logarithm at the point $(x, \log_b(x))$ equals $1/(x \ln(b))$.

The derivative of $\ln(x)$ is $1/x$; this implies that $\ln(x)$ is the unique antiderivative of $1/x$ that has the value 0 for $x = 1$. It is this very simple formula that motivated to qualify as "natural" the natural logarithm; this is also one of the main reasons of the importance of the constant e.

The derivative with a generalized functional argument $f(x)$ is

$$\frac{d}{dx}\ln f(x) = \frac{f'(x)}{f(x)}.$$

The quotient at the right hand side is called the logarithmic derivative of f. Computing $f'(x)$ by means of the derivative of $\ln(f(x))$ is known as logarithmic differentiation.[38] The antiderivative of the natural logarithm $\ln(x)$ is:[39]

$$\int \ln(x)\,dx = x\ln(x) - x + C.$$

Related formulas, such as antiderivatives of logarithms to other bases can be derived from this equation using the change of bases.[40]

Integral representation of the natural logarithm

The natural logarithm of t can be defined as the definite integral:

$$\ln t = \int_1^t \frac{1}{x}\,dx.$$

This definition has the advantage that it does not rely on the exponential function or any trigonometric functions; the definition is in terms of an integral of a simple reciprocal. As an integral, $\ln(t)$ equals the area between the x-axis and the graph of the function $1/x$, ranging from $x = 1$ to $x = t$. This is a consequence of the fundamental theorem of calculus and the fact that the derivative of $\ln(x)$ is $1/x$. Product and power logarithm formulas can be derived from this definition.[41] For example, the product formula $\ln(tu) = \ln(t) + \ln(u)$ is deduced as:

$$\ln(tu) \quad ¿ \int_1^{tu} \frac{1}{x}\,dx$$

$$¿ \quad [2] ¿ \ln(t) + \int_1^u \frac{1}{w}\,dw$$

$$¿ \quad ¿$$

The equality (1) splits the integral into two parts, while the equality (2) is a change of variable ($w = x/t$). In the illustration below, the splitting corresponds to dividing the area into the yellow and blue parts. Rescaling the

left hand blue area vertically by the factor t and shrinking it by the same factor horizontally does not change its size. Moving it appropriately, the area fits the graph of the function $f(x) = 1/x$ again. Therefore, the left hand blue area, which is the integral of $f(x)$ from t to tu is the same as the integral from 1 to u. This justifies the equality (2) with a more geometric proof.



A visual proof of the product formula of the natural logarithm

The power formula $\ln(t^r) = r\ln(t)$ may be derived in a similar way:

$$\begin{aligned}\ln\left(t^r\right) &= \int_1^{t^r} \frac{1}{x}\,dx \\ &= \int_1^{t} r\,\frac{1}{w}\,dw \\ &= r\ln(t)\end{aligned}$$

The second equality uses a change of variables (integration by substitution), $w = x^{1/r}$.

The sum over the reciprocals of natural numbers,

$$1+\frac{1}{2}+\frac{1}{3}+\cdots+\frac{1}{n}=\sum_{k=1}^{n}\frac{1}{k},$$

is called the harmonic series. It is closely tied to the natural logarithm: as n tends to infinity, the difference,

$$\sum_{k=1}^{n}\frac{1}{k}-\ln(n),$$

converges (i.e. gets arbitrarily close) to a number known as the Euler–Mascheroni constant $\gamma = 0.5772\ldots$. This relation aids in analyzing the performance of algorithms such as quicksort.[42]

Transcendence of the logarithm

Real numbers that are not algebraic are called transcendental;[43] for example, π and e are such numbers, but $\sqrt{2-\sqrt{3}}$ is not. Almost all real numbers are transcendental. The logarithm is an example of a transcendental function. The Gelfond–Schneider theorem asserts that logarithms usually take transcendental, i.e. "difficult" values.[44]

Calculation

Logarithms are easy to compute in some cases, such as $\log_{10}(1000) = 3$. In general, logarithms can be calculated using power series or the arithmetic–geometric mean, or be retrieved from a precalculated logarithm table that provides a fixed precision.[45][46] Newton's method, an iterative method to solve equations approximately, can also be used to calculate the logarithm, because its inverse function, the exponential function, can be computed efficiently.[47] Using look-up tables, CORDIC-like methods can be used to compute logarithms by using only the operations of addition and bit shifts.[48][49] Moreover, the binary logarithm algorithm calculates lb(x) recursively, based on repeated squarings of x, taking advantage of the relation

$$\log_2\left(x^2\right)=2\log_2 \vee x \vee .$$

Power series

Taylor series

For any real number z that satisfies $0 < z \leq 2$, the following formula holds:[nb 4][50]

$$\ln(z) \quad ¿\frac{\left(z-1¿^1\right.}{1}-\frac{\left(z-1¿^2\right.}{2}+\frac{\left(z-1¿^3\right.}{3}-\frac{\left(z-1¿^4\right.}{4}+\cdots$$
$$¿ \qquad\qquad\qquad\qquad\qquad\qquad ¿$$

Equating the function ln(z) to this infinite sum (series) is shorthand for saying that the function can be approximated to a more and more accurate value by the following expressions (known as partial sums):

$$(z-1), (z-1)-\frac{\left(z-1¿^2\right.}{2}, (z-1)-\frac{\left(z-1¿^2\right.}{2}+\frac{\left(z-1¿^3\right.}{3}, \ldots$$

For example, with $z = 1.5$ the third approximation yields 0.4167, which is about 0.011 greater than ln(1.5) = 0.405465, and the ninth approximation yields 0.40553, which is only about 0.0001 greater. The nth partial sum can approximate ln(z) with arbitrary precision, provided the number of summands n is large enough.

In elementary calculus, the series is said to converge to the function ln(z), and the function is the limit of the series. It is the Taylor series of the natural logarithm at $z = 1$. The Taylor series of ln(z) provides a particularly useful

approximation to ln(1 + z) when z is small, $|z| < 1$, since then

$$\ln(1+z) = z - \frac{z^2}{2} + \frac{z^3}{3} - \cdots \approx z.$$

For example, with $z = 0.1$ the first-order approximation gives ln(1.1) ≈ 0.1, which is less than 5% off the correct value 0.0953.

Inverse hyperbolic tangent

Another series is based on the inverse hyperbolic tangent function:

$$\ln(z) = 2 \cdot \operatorname{artanh}\frac{z-1}{z+1} = 2\left(\frac{z-1}{z+1} + \frac{1}{3}\left(\frac{z-1}{z+1}\right)^3 + \frac{1}{5}\left(\frac{z-1}{z+1}\right)^5 + \cdots\right),$$

for any real number $z > 0$.[nb 5][50] Using sigma notation, this is also written as

$$\ln(z) = 2\sum_{k=0}^{\infty} \frac{1}{2k+1}\left(\frac{z-1}{z+1}\right)^{2k+1}.$$

This series can be derived from the above Taylor series. It converges quicker than the Taylor series, especially if z is close to 1. For example, for $z = 1.5$, the first three terms of the second series approximate ln(1.5) with an error of about 3×10^{-6}. The quick convergence for z close to 1 can be taken advantage of in the following way: given a low-accuracy approximation $y \approx \ln(z)$ and putting

$$A = \frac{z}{\exp(y)},$$

the logarithm of z is:

$$\ln(z) = y + \ln(A).$$

The better the initial approximation y is, the closer A is to 1, so its logarithm can be calculated efficiently. A can be calculated using the exponential series, which converges quickly provided y is not too large. Calculating the logarithm of larger z can be reduced to smaller values of z by writing $z = a \cdot 10^b$, so that $\ln(z) = \ln(a) + b \cdot \ln(10)$.

A closely related method can be used to compute the logarithm of integers. Putting $z = \frac{n+1}{n}$ in the above series, it follows that:

$$\ln(n+1) = \ln(n) + 2\sum_{k=0}^{\infty} \frac{1}{2k+1}\left(\frac{1}{2n+1}\right)^{2k+1}.$$

If the logarithm of a large integer n is known, then this series yields a fast converging series for $\log(n+1)$, with a rate of convergence of $\left(\frac{1}{2n+1}\right)^2$.

Arithmetic–geometric mean approximation

The arithmetic–geometric mean yields high-precision approximations of the natural logarithm. Sasaki and Kanada showed in 1982 that it was particularly fast for precisions between 400 and 1000 decimal places, while Taylor series methods were typically faster when less precision was needed. In their work $\ln(x)$ is approximated to a precision of 2^{-p} (or p precise bits) by the following formula (due to Carl Friedrich Gauss):[51][52]

$$\ln(x) \approx \frac{\pi}{2M\left(1, 2^{2-m}/x\right)} - m\ln(2).$$

Here $M(x, y)$ denotes the arithmetic–geometric mean of x and y. It is obtained by repeatedly calculating the average $(x + y)/2$ (arithmetic mean) and $\sqrt{xy}$ (geometric mean) of x and y then let those two numbers become the next x and y. The two numbers quickly converge to a common limit which is the value of $M(x, y)$. m is chosen such that

$$x2^m > 2^{p/2}.$$

to ensure the required precision. A larger m makes the $M(x, y)$ calculation take more steps (the initial x and y are farther apart so it takes more steps to converge) but gives more precision. The constants π and $\ln(2)$ can be calculated with quickly converging series.

Feynman's algorithm

While at Los Alamos National Laboratory working on the Manhattan Project, Richard Feynman developed a bit-processing algorithm to compute the logarithm that is similar to long division and was later used in the Connection Machine. The algorithm relies on the fact that every real number x where $1 < x < 2$ can be represented as a product of distinct factors of the form $1 + 2^{-k}$. The algorithm sequentially builds that product P, starting with $P = 1$ and $k = 1$: if $P \cdot (1 + 2^{-k}) < x$, then it changes P to $P \cdot (1 + 2^{-k})$. It then increases k by one regardless. The algorithm stops when k is large enough to give the desired accuracy. Because $\log(x)$ is the sum of the terms of the form $\log(1 + 2^{-k})$ corresponding to those k for which the factor $1 + 2^{-k}$ was included in the product P, $\log(x)$ may be computed by simple

addition, using a table of $\log(1 + 2^{-k})$ for all k. Any base may be used for the logarithm table.[53]

Applications

Logarithms have many applications inside and outside mathematics. Some of these occurrences are related to the notion of scale invariance. For example, each chamber of the shell of a nautilus is an approximate copy of the next one, scaled by a constant factor. This gives rise to a logarithmic spiral.[54] Benford's law on the distribution of leading digits can also be explained by scale invariance.[55] Logarithms are also linked to self-similarity. For example, logarithms appear in the analysis of algorithms that solve a problem by dividing it into two similar smaller problems and patching their solutions.[56] The dimensions of self-similar geometric shapes, that is, shapes whose parts resemble the overall picture are also based on logarithms. Logarithmic scales are useful for quantifying the relative change of a value as opposed to its absolute difference. Moreover, because the logarithmic function $\log(x)$ grows very slowly for large x, logarithmic scales are used to compress large-scale scientific data. Logarithms also occur in numerous scientific formulas, such as the Tsiolkovsky rocket equation, the Fenske equation, or the Nernst equation.

Logarithmic scale

Scientific quantities are often expressed as logarithms of other quantities, using a *logarithmic scale*. For example, the decibel is a unit of measurement associated with logarithmic-scale quantities. It is based on the common logarithm of ratios—10 times the common logarithm of a power ratio or 20 times the common logarithm of a voltage ratio. It is used to quantify the attenuation or amplification of electrical signals,[57] to describe power levels of sounds in acoustics,[58] and the absorbance of light in the fields of spectrometry and optics. The signal-to-noise ratio describing the amount of unwanted noise in relation to a (meaningful) signal is also measured in decibels.[59] In a similar vein, the peak signal-to-noise ratio is commonly used to assess the quality of sound and image compression methods using the logarithm.[60]

The strength of an earthquake is measured by taking the common logarithm of the energy emitted at the quake. This is used in the moment magnitude scale or the Richter magnitude scale. For example, a 5.0 earthquake releases

32 times ($10^{1.5}$) and a 6.0 releases 1000 times (10^3) the energy of a 4.0.[61] Apparent magnitude measures the brightness of stars logarithmically.[62] In chemistry the negative of the decimal logarithm, the decimal **cologarithm**, is indicated by the letter p.[63] For instance, pH is the decimal cologarithm of the activity of hydronium ions (the form hydrogen ions H^+ take in water).[64] The activity of hydronium ions in neutral water is 10^{-7} $mol \cdot L^{-1}$, hence a pH of 7. Vinegar typically has a pH of about 3. The difference of 4 corresponds to a ratio of 10^4 of the activity, that is, vinegar's hydronium ion activity is about 10^{-3} $mol \cdot L^{-1}$.

Semilog (log–linear) graphs use the logarithmic scale concept for visualization: one axis, typically the vertical one, is scaled logarithmically. For example, the chart at the right compresses the steep increase from 1 million to 1 trillion to the same space (on the vertical axis) as the increase from 1 to 1 million. In such graphs, exponential functions of the form $f(x) = a \cdot b^x$ appear as straight lines with slope equal to the logarithm of b. Log-log graphs scale both axes logarithmically, which causes functions of the form $f(x) = a \cdot x^k$ to be depicted as straight lines with slope equal to the exponent k. This is applied in visualizing and analyzing power laws.[65]

Psychology

Logarithms occur in several laws describing human perception:[66][67] Hick's law proposes a logarithmic relation between the time individuals take to choose an alternative and the number of choices they have.[68] Fitts's law predicts that the time required to rapidly move to a target area is a logarithmic function of the ratio between the distance to a target and the size of the target.[69] In psychophysics, the Weber–Fechner law proposes a logarithmic relationship between stimulus and sensation such as the actual vs. the perceived weight of an item a person is carrying.[70] (This "law", however, is less realistic than more recent models, such as Stevens's power law.[71])

Psychological studies found that individuals with little mathematics education tend to estimate quantities logarithmically, that is, they position a number on an unmarked line according to its logarithm, so that 10 is positioned as close to 100 as 100 is to 1000. Increasing education shifts this to a linear estimate (positioning 1000 10 times as far away) in some circumstances, while logarithms are used when the numbers to be plotted are difficult to plot linearly.[72][73]

Probability theory and statistics

Logarithms arise in probability theory: the law of large numbers dictates that, for a fair coin, as the number of coin-tosses increases to infinity, the observed proportion of heads approaches one-half. The fluctuations of this proportion about one-half are described by the law of the iterated logarithm.[74]

Logarithms also occur in log-normal distributions. When the logarithm of a random variable has a normal distribution, the variable is said to have a log-normal distribution.[75] Log-normal distributions are encountered in many fields, wherever a variable is formed as the product of many independent positive random variables, for example in the study of turbulence.[76]

Logarithms are used for maximum-likelihood estimation of parametric statistical models. For such a model, the likelihood function depends on at least one parameter that must be estimated. A maximum of the likelihood function occurs at the same parameter-value as a maximum of the logarithm of the likelihood (the "*log likelihood*"), because the logarithm is an increasing function. The log-likelihood is easier to maximize, especially for the multiplied likelihoods for independent random variables.[77]

Benford's law describes the occurrence of digits in many data sets, such as heights of buildings. According to Benford's law, the probability that the first decimal-digit of an item in the data sample is d (from 1 to 9) equals $\log_{10}(d+1) - \log_{10}(d)$, *regardless* of the unit of measurement.[78] Thus, about 30% of the data can be expected to have 1 as first digit, 18% start with 2, etc. Auditors examine deviations from Benford's law to detect fraudulent accounting.[79]

The logarithm transformation is a type of data transformation used to bring the empirical distribution closer to the assumed one.

Computational complexity

Analysis of algorithms is a branch of computer science that studies the performance of algorithms (computer programs solving a certain problem).[80] Logarithms are valuable for describing algorithms that divide a problem into smaller ones, and join the solutions of the subproblems.[81]

For example, to find a number in a sorted list, the binary search algorithm checks the middle entry and proceeds with the half before or after the middle entry if the number is still not found. This algorithm requires, on

average, $\log_2(N)$ comparisons, where N is the list's length.[82] Similarly, the merge sort algorithm sorts an unsorted list by dividing the list into halves and sorting these first before merging the results. Merge sort algorithms typically require a time approximately proportional to $N \cdot \log(N)$.[83] The base of the logarithm is not specified here, because the result only changes by a constant factor when another base is used. A constant factor is usually disregarded in the analysis of algorithms under the standard uniform cost model.[84]

A function $f(x)$ is said to grow logarithmically if $f(x)$ is (exactly or approximately) proportional to the logarithm of x. (Biological descriptions of organism growth, however, use this term for an exponential function.[85]) For example, any natural number N can be represented in binary form in no more than $\log_2 N + 1$ bits. In other words, the amount of memory needed to store N grows logarithmically with N.

Entropy and chaos

Entropy is broadly a measure of the disorder of some system. In statistical thermodynamics, the entropy S of some physical system is defined as

$$S = -k \sum_i p_i \ln(p_i).$$

The sum is over all possible states i of the system in question, such as the positions of gas particles in a container. Moreover, p_i is the probability that the state i is attained and k is the Boltzmann constant. Similarly, entropy in information theory measures the quantity of information. If a message recipient may expect any one of N possible messages with equal likelihood, then the amount of information conveyed by any one such message is quantified as $\log_2 N$ bits.[86]

Lyapunov exponents use logarithms to gauge the degree of chaoticity of a dynamical system. For example, for a particle moving on an oval billiard table, even small changes of the initial conditions result in very different paths of the particle. Such systems are chaotic in a deterministic way, because small measurement errors of the initial state predictably lead to largely different final states.[87] At least one Lyapunov exponent of a deterministically chaotic system is positive.

Fractals

Logarithms occur in definitions of the dimension of fractals.[88] Fractals are geometric objects that are self-similar in the sense that small parts reproduce, at least roughly, the entire global structure. The Sierpinski triangle (pictured) can be covered by three copies of itself, each having sides half the original length. This makes the Hausdorff dimension of this structure ln(3)/ln(2) ≈ 1.58. Another logarithm-based notion of dimension is obtained by counting the number of boxes needed to cover the fractal in question.

Music

Logarithms are related to musical tones and intervals. In equal temperament tunings, the frequency ratio depends only on the interval between two tones, not on the specific frequency, or pitch, of the individual tones. In the 12-tone equal temperament tuning common in modern Western music, each octave (doubling of frequency) is broken into twelve equally spaced intervals called semitones. For example, if the note *A* has a frequency of 440 Hz then the note *B-flat* has a frequency of 466 Hz. The interval between *A* and *B-flat* is a semitone, as is the one between *B-flat* and *B* (frequency 493 Hz). Accordingly, the frequency ratios agree:

$$\frac{466}{440} \approx \frac{493}{466} \approx 1.059 \approx \sqrt[12]{2}.$$

Intervals between arbitrary pitches can be measured in octaves by taking the base-2 logarithm of the frequency ratio, can be measured in equally tempered semitones by taking the base-$2^{1/12}$ logarithm (12 times the base-2 logarithm), or can be measured in cents, hundredths of a semitone, by taking the base-$2^{1/1200}$ logarithm (1200 times the base-2 logarithm). The latter is used for finer encoding, as it is needed for finer measurements or non-equal temperaments.[89]

Interval (the two tones are played at the same time)	1/12 tone play	Semitone play	Just major third play	Major third play	Tritone play	Octave play
Frequency ratio r	$2^{\frac{1}{72}} \approx 1.00$	$2^{\frac{1}{12}} \approx 1.05$	$\frac{5}{4} = 1.25$	$2^{\frac{4}{12}}$ ¿ ¿ $\sqrt[3]{2}$ ¿	$2^{\frac{6}{12}}$ ¿ ¿ $\sqrt{2}$ ¿	$2^{\frac{12}{12}} = 2$

Number of semitones $12\log_2 r$	$\frac{1}{6}$	1	≈ 3.8631	4	6	12
Number of cents $1200\log_2 r$	$16\frac{2}{3}$	100	≈ 386.31	400	600	1200

Number theory

Natural logarithms are closely linked to counting prime numbers (2, 3, 5, 7, 11, ...), an important topic in number theory. For any integer x, the quantity of prime numbers less than or equal to x is denoted $\pi(x)$. The prime number theorem asserts that $\pi(x)$ is approximately given by

$$\frac{x}{\ln(x)},$$

in the sense that the ratio of $\pi(x)$ and that fraction approaches 1 when x tends to infinity.[90] As a consequence, the probability that a randomly chosen number between 1 and x is prime is inversely proportional to the number of decimal digits of x. A far better estimate of $\pi(x)$ is given by the offset logarithmic integral function Li(x), defined by

$$Li(x) = \int_2^x \frac{1}{\ln(t)}\,dt.$$

The Riemann hypothesis, one of the oldest open mathematical conjectures, can be stated in terms of comparing $\pi(x)$ and Li(x).[91] The Erdős–Kac theorem describing the number of distinct prime factors also involves the natural logarithm.

The logarithm of n factorial, $n! = 1 \cdot 2 \cdot \ldots \cdot n$, is given by

$$\ln(n!) = \ln(1) + \ln(2) + \cdots + \ln(n).$$

This can be used to obtain Stirling's formula, an approximation of $n!$ for large n.[92]

Generalizations

Complex logarithm

All the complex numbers a that solve the equation

$$e^a = z$$

are called *complex logarithms* of z, when z is (considered as) a complex number. A complex number is commonly represented as $z = x + iy$, where x and y are real numbers and i is an imaginary unit, the square of which is -1. Such a number can be visualized by a point in the complex plane, as shown at the right. The polar form encodes a non-zero complex number z by its absolute value, that is, the (positive, real) distance r to the origin, and an angle between the real (x) axis Re and the line passing through both the origin and z. This angle is called the argument of z.

The absolute value r of z is given by

$$r = \sqrt{x^2 + y^2}.$$

Using the geometrical interpretation of sine and cosine and their periodicity in 2π, any complex number z may be denoted as

$$\begin{aligned} z &= x + iy \\ &= r(\cos(\varphi + 2k\pi) + i\sin(\varphi + 2k\pi)), \end{aligned}$$

for any integer number k. Evidently the argument of z is not uniquely specified: both φ and $\varphi' = \varphi + 2k\pi$ are valid arguments of z for all integers k, because adding $2k\pi$ radians or $k \cdot 360°$[nb 6] to φ corresponds to "winding" around the origin counter-clock-wise by k turns. The resulting complex number is always z, as illustrated at the right for $k = 1$. One may select exactly one of the possible arguments of z as the so-called *principal argument*, denoted Arg(z), with a capital A, by requiring φ to belong to one, conveniently selected turn, e.g. $-\pi < \varphi \leq \pi$[93] or $0 \leq \varphi < 2\pi$.[94] These regions, where the argument of z is uniquely determined are called *branches* of the argument function.

Euler's formula connects the trigonometric functions sine and cosine to the complex exponential:

$$e^{i\varphi} = \cos\varphi + i\sin\varphi.$$

Using this formula, and again the periodicity, the following identities hold:[95]

$$\begin{aligned} z &= r(\cos\varphi + i\sin\varphi) \\ &= re^{i(\varphi + 2k\pi)} \\ &= e^{\ln(r) + i(\varphi + 2k\pi)} = e^{a_k}, \end{aligned}$$

where ln(r) is the unique real natural logarithm, a_k denote the complex logarithms of z, and k is an arbitrary integer. Therefore, the complex logarithms of z, which are all those complex values a_k for which the a_k-th power of e equals z, are the infinitely many values

$$a_k = \ln(r) + i(\varphi + 2k\pi),$$

for arbitrary integers k.

Taking k such that $\varphi + 2k\pi$ is within the defined interval for the principal arguments, then a_k is called the *principal value* of the logarithm, denoted Log(z), again with a capital L. The principal argument of any positive real number x is 0; hence Log(x) is a real number and equals the real (natural) logarithm. However, the above formulas for logarithms of products and powers do *not* generalize to the principal value of the complex logarithm.[96]

The illustration at the right depicts Log(z), confining the arguments of z to the interval $(-\pi, \pi]$. This way the corresponding branch of the complex logarithm has discontinuities all along the negative real x axis, which can be seen in the jump in the hue there. This discontinuity arises from jumping to the other boundary in the same branch, when crossing a boundary, i.e. not changing to the corresponding k-value of the continuously neighboring branch. Such a locus is called a branch cut. Dropping the range restrictions on the argument makes the relations "argument of z", and consequently the "logarithm of z", multi-valued functions.

Inverses of other exponential functions

Exponentiation occurs in many areas of mathematics and its inverse function is often referred to as the logarithm. For example, the logarithm of a matrix is the (multi-valued) inverse function of the matrix exponential.[97] Another example is the p-adic logarithm, the inverse function of the p-adic exponential. Both are defined via Taylor series analogous to the real case.[98] In the context of differential geometry, the exponential map maps the tangent space at a point of a manifold to a neighborhood of that point. Its inverse is also called the logarithmic (or log) map.[99]

In the context of finite groups exponentiation is given by repeatedly multiplying one group element b with itself. The discrete logarithm is the integer n solving the equation

$$b^n = x,$$

where x is an element of the group. Carrying out the exponentiation can be done efficiently, but the discrete logarithm is believed to be very hard to

calculate in some groups. This asymmetry has important applications in public key cryptography, such as for example in the Diffie–Hellman key exchange, a routine that allows secure exchanges of cryptographic keys over unsecured information channels.[100] Zech's logarithm is related to the discrete logarithm in the multiplicative group of non-zero elements of a finite field.[101]

Further logarithm-like inverse functions include the *double logarithm* $\ln(\ln(x))$, the *super- or hyper-4-logarithm* (a slight variation of which is called iterated logarithm in computer science), the Lambert W function, and the logit. They are the inverse functions of the double exponential function, tetration, of $f(w) = we^w$,[102] and of the logistic function, respectively.[103]

Related concepts

From the perspective of group theory, the identity $\log(cd) = \log(c) + \log(d)$ expresses a group isomorphism between positive reals under multiplication and reals under addition. Logarithmic functions are the only continuous isomorphisms between these groups.[104] By means of that isomorphism, the Haar measure (Lebesgue measure) dx on the reals corresponds to the Haar measure dx/x on the positive reals.[105] The non-negative reals not only have a multiplication, but also have addition, and form a semiring, called the probability semiring; this is in fact a semifield. The logarithm then takes multiplication to addition (log multiplication), and takes addition to log addition (LogSumExp), giving an isomorphism of semirings between the probability semiring and the log semiring.

Logarithmic one-forms df/f appear in complex analysis and algebraic geometry as differential forms with logarithmic poles.[106]

The polylogarithm is the function defined by

$$Li_s(z) = \sum_{k=1}^{\infty} \frac{z^k}{k^s}.$$

It is related to the natural logarithm by $\mathrm{Li}_1(z) = -\ln(1 - z)$. Moreover, $\mathrm{Li}_s(1)$ equals the Riemann zeta function $\zeta(s)$.[107]

See also

- Decimal exponent (dex)
- Exponential function

- Index of logarithm articles

Read more (what I didn't insert back then)

Logarithm



Plots of logarithm functions, with three commonly used bases. The special points $\log_b b = 1$ are indicated by dotted lines, and all curves intersect in $\log_b 1 = 0$.

Motivation



The graph of the logarithm base 2 crosses the *x*-axis at $x = 1$ and passes through the points (2, 1), (4, 2), and (8, 3), depicting, e.g., $\log_2(8) = 3$ and $2^3 = 8$. The graph gets arbitrarily close to the *y*-axis, but does not meet it.

Particular bases



Overlaid graphs of the logarithm for bases $\frac{1}{2}$, 2, and *e*

Logarithm tables, slide rules, and historical applications



LOGARITHMS, (from λογος *ratio*, and αριθμος *number*), the indices of the ratios of numbers to one another; being a ſeries of numbers in arithmetical progreſſion, correſponding to others in geometrical progreſſion; by means of which, arithmetical calculations can be made with much more eaſe and expedition than otherwiſe.

The 1797 *Encyclopædia Britannica* explanation of logarithms

Analytic properties

Graph of the logarithm function



The graph of the logarithm function $\log_b(x)$ (blue) is obtained by reflecting the graph of the function b^x (red) at the diagonal line ($x = y$).

Derivative and antiderivative



The graph of the natural logarithm (green) and its tangent at $x = 1.5$ (black)

Integral representation of the natural logarithm



The natural logarithm of t is the shaded area underneath the graph of the function $f(x) = 1/x$.

Calculation

The logarithm keys (LOG for base 10 and LN for base e) on a TI-83 Plus graphing calculator

Power series

Taylor series



The Taylor series of ln(*z*) centered at *z* = 1. The animation shows the first 10 approximations along with the 99th and 100th. The approximations do not converge beyond a distance of 1 from the center.

Applications

A nautilus shell displaying a logarithmic spiral

Logarithmic scale

A logarithmic chart depicting the value of one Goldmark in Papiermarks during the German hyperinflation in the 1920s

Probability theory and statistics



Three probability density functions (PDF) of random variables with log-normal distributions. The location parameter μ, which is zero for all three of the PDFs shown, is the mean of the logarithm of the random variable, not the mean of the variable itself.



Distribution of first digits (in %, red bars) in the population of the 237 countries of the world. Black dots indicate the distribution predicted by Benford's law.

Entropy and chaos



Billiards on an oval billiard table. Two particles, starting at the center with an angle differing by one degree, take paths that diverge chaotically because of reflections at the boundary.

Fractals

The Sierpinski triangle (at the right) is constructed by repeatedly replacing equilateral triangles by three smaller ones.

Music



Four different octaves shown on a linear scale, then shown on a logarithmic scale (as the ear hears them)

Generalizations

Complex logarithm



Polar form of $z = x + iy$.
Both φ and φ' are arguments of z.



The principal branch ($-\pi$, π) of the complex logarithm, Log(z). The black point at $z =$

1 corresponds to absolute value zero and brighter colors refer to bigger absolute values. The hue of the color encodes the argument of Log(z).